

New antimicrobial agents for the treatment of Gram-positive bacterial infections

D. Y. Aksoy¹ and S. Unal²

¹Department of Internal Medicine and ²Department of Internal Medicine, Section of Infectious Diseases, Medical Faculty, Hacettepe University, Ankara, Turkey

ABSTRACT

Since the 1970s, resistance to antimicrobial agents has become an escalating problem. In the last 25 years, treatment of infections caused by Gram-positive bacteria has been more problematical than ever, with infections being caused by multidrug-resistant organisms, particularly methicillin-resistant staphylococci, penicillin- and erythromycin-resistant pneumococci, and vancomycin-resistant enterococci. There is a continuing effort in the pharmaceutical industry to develop new antimicrobial agents for the treatment of resistant infections. Linezolid, quinupristin–dalfopristin, daptomycin, tigecycline, new glycopeptides and ceftobiprole are the main agents recently introduced or under clinical development. This review summarises their major properties, the results of recent studies with these agents, and future treatment possibilities.

Keywords Ceftobiprole, daptomycin, linezolid, quinupristin–dalfopristin, review, tigecycline

Accepted: 5 November 2007

Clin Microbiol Infect 2008; **14**: 411–420

INTRODUCTION

The introduction of benzylpenicillin in the 1940s ushered in an era of effective treatment of bacterial infections, but it soon became apparent that some strains of *Staphylococcus aureus* were drug-resistant because of the production of β -lactamase. Since the mid-1970s, resistance to antimicrobial agents has become an escalating problem [1]. A striking change during the past quarter-century has been the increasing importance of infections caused by Gram-positive bacteria, which might have resulted from the previous emphasis placed on controlling Gram-negative bacterial infections. Today, it is necessary to deal with infections caused by multidrug-resistant organisms, particularly methicillin-resistant staphylococci, penicillin- and erythromycin-resistant pneumococci, and vancomycin-resistant enterococci (VRE). The emergence and rapid spread of strains of methicillin-resistant *Staph. aureus* (MRSA), which are resistant not only

to all β -lactams, but also to the other main antibiotic classes, has resulted in an increased use of glycopeptide antibiotics, i.e., vancomycin and teicoplanin. Unfortunately, VRE were detected in 1986, both in France and the UK, and multidrug-resistant strains of enterococci are now encountered throughout Europe and the USA [2–4]. Within 1 year of their initial detection, resistant strains were associated with infections in the USA, and they accounted for 26% of isolates of enterococci from blood in the USA by the year 2000. Since 1996, vancomycin-intermediate *Staph. aureus* (VISA) isolates with a vancomycin MIC of 8–16 mg/L have been reported, and this has been followed since 2002 by reports of vancomycin-resistant *Staph. aureus* (MIC ≥ 32 mg/L) [5–8].

As a result of these developments, there is an urgent need for effective new antimicrobial agents, as well as for prudent use of existing agents. Linezolid, quinupristin–dalfopristin, daptomycin, tigecycline, new glycopeptides and ceftobiprole are the main new agents that have recently become available for use or are under clinical development. This review summarises their properties, the results of recent studies with these agents, and future treatment possibilities.

Corresponding author and reprint requests: D. Y. Aksoy, Angora Evleri E-2/31, 06530 Beysukent, Ankara, Turkey
E-mail: duyguayaks@yahoo.com

LINEZOLID

Linezolid (Zyvox[®]) is the first of a new class of antimicrobial agents, the oxazolidinones. Linezolid is a synthetic antibiotic that, depending on the organism and the linezolid concentration, has a bacteriostatic or bactericidal effect by inhibiting protein synthesis at the ribosomal level and by preventing the formation of the protein initiation complex [9]. Linezolid is active against most Gram-positive bacteria, including methicillin-sensitive *Staph. aureus* (MSSA) and MRSA, *Streptococcus pneumoniae* (including multidrug-resistant strains), *Streptococcus pyogenes* and *Streptococcus agalactiae*. Rare occurrences of linezolid resistance among VRE in North America (0.8–1.8%), caused by a G2576U ribosomal mutation, have been reported, and an intrinsic resistance gene rendering a clinical strain of MRSA resistant to linezolid has been described [10–13]. Although resistant strains of *Staph. aureus* have been reported, linezolid remains highly active against MRSA (MIC₉₀ 2 mg/L) [12–15].

Linezolid was approved by the US Food and Drug Administration (FDA) in 2000 for the treatment of uncomplicated and complicated skin and soft-tissue infections, including diabetic foot infections without concomitant osteomyelitis, community-acquired and nosocomial pneumonia, and vancomycin-resistant *Enterococcus faecium* infections, including cases with concurrent bacteraemia [9]. Activity against *Nocardia* spp. and *Mycobacterium* spp. has also been demonstrated, including cases of central nervous system infection and infective endocarditis [14–18].

The high bioavailability of linezolid, whether administered orally or intravenously, with a standard twice-daily dose of 600 mg, makes it a suitable alternative for the treatment of infections that require the prolonged use of antimicrobial agents [19,20]. Tissue penetration is good, with marked accumulation in sweat, saliva and epithelial lining. Linezolid is reported to be equally effective or even superior to vancomycin for the treatment of pneumonia and soft-tissue infections [21–23]. However, recent meta-analyses have suggested that linezolid's apparent superiority for the treatment of pneumonia and bone and joint infections has limitations. Cases of MRSA endocarditis that failed to respond to linezolid treatment have also been reported [15,24,25].

Linezolid is generally well-tolerated, with the most common adverse effects being diarrhoea, headache and nausea. Like many other antibiotics, it may predispose to pseudo-membranous colitis following the overgrowth of *Clostridium difficile* [26]. Myelosuppression (including anaemia, leukopenia, pancytopenia, and thrombocytopenia) has been reported in patients taking linezolid. In cases with a known treatment outcome, the affected haematological parameters rose to pre-treatment levels when linezolid was discontinued. Complete blood counts should be monitored weekly in patients taking linezolid, and particularly in patients who are prescribed linezolid for >2 weeks, patients with pre-existing myelosuppression, patients receiving concomitant drugs that produce bone marrow suppression, and patients with a chronic infection who have received previous or concomitant antibiotic therapy. Discontinuation of linezolid therapy should be considered for patients who develop, or have worsening, myelosuppression. Spontaneous reports of serotonin syndrome associated with the co-administration of linezolid and serotonergic agents, including such anti-depressants as selective serotonin re-uptake inhibitors, have been reported. Linezolid has not been studied in patients with uncontrolled hypertension, pheochromocytoma, carcinoid syndrome or untreated hyperthyroidism [27].

Lactic acidosis, peripheral neuropathy and optic neuropathy are other adverse effects of linezolid treatment. A recent study that compared linezolid to vancomycin, oxacillin and dicloxacillin (comparator antibiotics) for the treatment of seriously-ill patients with intravascular catheter-related bloodstream infections, including patients with catheter-site infections, reported a higher mortality rate in the linezolid arm. There was no difference in the mortality rate of patients with Gram-positive infections who were treated with other antibiotics. Furthermore, the mortality rate was higher only for patients who were infected only with Gram-negative organisms, or who were infected with both Gram-positive and Gram-negative organisms, or who had no infection when they entered the study and were treated with linezolid. Therefore, if infection with Gram-negative bacteria is known or suspected, appropriate therapy should be started immediately (<http://www.fda.gov/cder/drug/InfoSheets/HCP/linezolidHCP.htm>).

Linezolid is an effective alternative antimicrobial agent for the treatment of Gram-positive infections, but because of the absence of placebo-controlled double-blind studies, linezolid should be used with extreme caution in patients with vancomycin-induced nephrotoxicity or a documented absence of response to vancomycin.

QUINUPRISTIN-DALFOPRISTIN

Quinupristin-dalfopristin (Synercid[®]) is a fixed mixture (30:70) of semi-synthetic streptogramin derivatives. These compounds enter bacterial cells by diffusion and bind to different sites on the 50S ribosomal subunit, resulting in an irreversible inhibition of bacterial protein synthesis [28]. Dalfopristin blocks the reaction catalysed by the peptidyl transferase catalytic centre of the 50S ribosome by inhibiting substrate attachment to the P-site and A-site of the ribosome. Quinupristin inhibits peptide chain elongation. The synergic effect of this drug combination appears to result from the fact that it targets both early and late stages of protein synthesis [29]. Thus, quinupristin-dalfopristin is bacteriostatic against *E. faecium* and bactericidal against MSSA, MRSA and *Strep. pyogenes*. It is ineffective against *Enterococcus faecalis*. In the USA, the only FDA-approved use of quinupristin-dalfopristin as an anti-staphylococcal agent is for the treatment of adults with complicated skin and skin-structure infections caused by MSSA. Otherwise, the drug is approved for complicated skin and skin-structure infections caused by *Strep. pyogenes*, and for serious VRE infections associated with bacteraemia [30]. Resistance in *E. faecium* was reported by the SENTRY antimicrobial surveillance programme in Europe and North America to be 10.0% and 0.6%, respectively [10].

The results of studies in which the effectiveness of quinupristin-dalfopristin for the treatment of skin and soft-tissue infections was compared to that of other agents have generally been satisfactory, despite the fact that quinupristin-dalfopristin is inferior to other agents for the treatment of pneumonia and infective endocarditis [24,31].

Quinupristin-dalfopristin can only be administered by the intravenous route in a dextrose 5% w/v solution. Drug elimination is through bile into faeces, but clearance may be slightly impaired in patients with renal insufficiency. The drug has significant toxicity problems, including

arthralgia-myalgia syndrome and venous intolerance. Pain and inflammation at the infusion site is experienced by up to 74% of patients [31-33]. Hyper-bilirubinaemia and liver toxicity can also occur. Quinupristin-dalfopristin has been shown to be a major inhibitor of the activity of the cytochrome P450 3A4 isoenzyme, and drug interactions (especially with cyclosporine) should be monitored carefully during therapy. Although the drug itself does not induce QTc prolongation, it can interfere with the metabolism of other drug products that are associated with QTc prolongation [33,34].

Quinupristin-dalfopristin is the first parenteral streptogramin and offers a unique alternative treatment for infections caused by multidrug-resistant Gram-positive bacteria, but the broad spectrum of adverse effects makes it an inferior choice to other agents.

DAPTOMYCIN

Daptomycin (Cubicin[®]) is the first of a new class of cyclic lipopeptides. This agent is derived from the fermentation of a strain of *Streptomyces roseosporus*. Originally developed during the early 1980s, daptomycin was initially shelved because of concerns about skeletal-muscle toxicity. This effect was not seen with lower doses, and daptomycin received approval by the FDA in September 2003 for the treatment of complicated skin and soft-tissue infections.

Daptomycin attaches to the cytoplasmic membrane of Gram-positive bacteria, forming a channel that allows the efflux of potassium, causing depolarisation of the membrane, along with dysfunction of macromolecular synthesis and collapse of the organism without lysis. This mechanism of action is concentration-dependent, free calcium ion-dependent, rapid and unique [35]. Some other bactericidal antibiotics, most notably β -lactams, cause bacterial cells to lyse. This is potentially harmful, as it may release bacterial endotoxins and other inflammatory cell components into the circulation, triggering cytokine cascades, and potentially leading to septic shock and multiple organ failure [36]. The antimicrobial effect of daptomycin is concentration-dependent, with the optimal dose for serious bloodstream and endovascular infections reported to be 6 mg/kg once-daily. The recommended dose for skin and skin-structure infections is 4 mg/kg once-daily.

Daptomycin also has a post-antibiotic effect lasting 1.5–6 h [37].

Although daptomycin is approved by the FDA for the treatment of complicated skin and skin-structure infections that involve MRSA and other Gram-positive bacteria, its rapid bactericidal effect makes it appropriate for the treatment of other kinds of infection. The bactericidal activity of daptomycin, vancomycin, linezolid and quinupristin–dalbapristin against MRSA and VISA has been compared using in-vitro time-kill studies [38]. Against all organisms tested, daptomycin had bactericidal activity equal to or greater than that of the other agents [39].

Daptomycin has efficacy comparable to standard therapy for the treatment of skin and skin-structure infections. It has been used successfully to treat bone and joint infections [40,41]. Successful results have also been reported for the treatment of bacteraemia and right-sided infective endocarditis, but not for the treatment of community-acquired pneumonia. When daptomycin was compared with standard therapy for bacteraemia and endocarditis caused by *Staph. aureus*, a successful outcome was documented in 53 of 120 of patients who received daptomycin, compared with 48 of 115 patients who received standard therapy (44.2% vs. 41.7%, respectively; absolute difference, 2.4%; 95% CI 10.2–15.1) [42–44]. The lack of efficacy of daptomycin in treating community-acquired pneumonia is thought to be a consequence of a reduction of daptomycin activity in the presence of lung surfactant [45].

Resistance to daptomycin is rare. Spontaneous resistance is uncommon, emerging *in vitro* at a rate of $<1 \times 10^{-10}$, but resistance can be induced by serial passage in increasing concentrations of the antibiotic [46]. There have been several reports of daptomycin resistance emerging in clinical isolates of MRSA from patients who received prolonged treatment [47,48]. The once-daily dosing schedule and the favourable safety profile (except for some concerns regarding rhabdomyolysis and neuropathy) make daptomycin an attractive option for the treatment of Gram-positive infections [44].

TIGECYCLINE

Tigecycline (Tygacil®) is a new, semi-synthetic glycylcycline, which is a new class of antimicrobial agent. Glycocides are derivatives of tetracyclines, and have broad-spectrum activity against

susceptible and multidrug-resistant strains of bacteria. Tigecycline, the first glycylcycline, demonstrates potent in-vitro activity against a wide range of Gram-positive and Gram-negative bacteria, including MRSA and tetracycline-resistant *Staph. aureus*, as well as many multidrug-resistant Gram-negative pathogens and anaerobes [49]. The potent in-vitro activity of tigecycline resulted in its approval by the FDA for the treatment of skin, soft-tissue and intra-abdominal infections [50].

Like tetracycline, tigecycline exerts its mechanism of action by reversibly binding to the 30S ribosomal subunit and inhibiting protein translation in bacteria. This blockade prevents the entry of aminoacyl tRNA molecules into the A-site of the ribosome, resulting in the loss of peptide formation [51,52]. Tigecycline is highly effective *in vivo* against most Gram-positive organisms, including *Staph. aureus*, coagulase-negative staphylococci, *Enterococcus* spp., *Strep. pneumoniae*, and group A, group B and viridans streptococci. It also has good in-vitro activity against most Gram-negative organisms and anaerobes. Interestingly, some atypical microorganisms, e.g., *Mycobacterium abscessus*, *Mycobacterium chelonae*, the *Mycobacterium fortuitum* group, *Mycobacterium marinum*, *Chlamydia pneumoniae*, *Mycoplasma hominis*, *Mycoplasma pneumonia* and *Ureaplasma urealyticum*, are also susceptible to tigecycline [53]. Tigecycline has demonstrated excellent overall tissue penetration in animal studies, including bone, bone marrow, salivary gland, thyroid, spleen, kidney and cerebrospinal fluid [53].

Phase 3 randomised double-blind studies have confirmed the efficacy of tigecycline for the treatment of skin and skin-structure infections, and for intra-abdominal infections. Sacchidanand *et al.* [54] reported that tigecycline was not inferior to a combination of aztreonam and vancomycin for the treatment of complicated skin and skin-structure infections. Cure rates and microbiological eradication rates were similar for tigecycline and a comparator group of antibiotics, at 82.9% and 82.3%, respectively [54]. In two other phase 3 double-blind studies involving 833 clinically evaluable patients (422 treated with tigecycline and 411 treated with vancomycin and aztreonam), clinical response rates were similar for tigecycline (86.5%) (95% CI 82.9–89.6%) and vancomycin–aztreonam (88.6%) (95% CI 85.1–91.5%) [55].

In a pooled analysis of two phase 3 double-blind trials designed to evaluate the safety and efficacy of tigecycline in comparison with that of imipenem–cilastatin, clinical cure rates for the microbiologically evaluable group among 1642 adults with complicated intra-abdominal infections were 86.1% for tigecycline and 86.2% for imipenem–cilastatin (95% CI for the difference –4.5 to 4.4%; $p < 0.0001$ for non-inferiority) [56]. Similar results were obtained in another study which demonstrated that tigecycline was not inferior for treating complicated intra-abdominal infections [57].

Clinically significant organ toxicity has not been observed in association with tigecycline use during clinical trials. Nausea and vomiting are the most common side-effects, and these are dose-limited and are not diminished by reducing the rate of drug infusion [58]. The low potential of the drug for the development of resistance, which is almost a rule for other tetracyclines, combined with its broad-spectrum activity against multidrug-resistant pathogens, provides key advantages. Therefore, tigecycline presents a genuinely new therapeutic option for the treatment of infections caused by multidrug-resistant bacteria.

NEW GLYCOPEPTIDES

Oritavancin, telavancin and dalbavancin are new glycopeptides currently in clinical development, and appear to be potent molecules with favourable pharmacokinetic and pharmacodynamic properties.

Oritavancin

This agent was obtained by reductive alkylation with 4'-chloro-biphenylcarboxaldehyde of chloroeremomycin, which differs from vancomycin by the addition of a 4-epi-vancosamine sugar, and the replacement of vancosamine of the disaccharide moiety by an epivancosamine [59]. Although oritavancin has a general spectrum of activity comparable to that of vancomycin, it offers considerable advantages in terms of intrinsic activity (especially against streptococci). Unlike vancomycin, oritavancin can dimerise, leading to a cooperative interaction with two stems of the growing peptidoglycan chain. The lipophilic side-chain assists in membrane-

anchoring by hydrophobic interactions, stabilising the dimer in the most favourable position. Evidence also exists for another mechanism of action of oritavancin, namely inhibition of the transglycosylation step of cell-wall synthesis [60,61].

Oritavancin is active against most isolates of streptococci and staphylococci, as well as *Peptostreptococcus* spp., *Propionibacterium acnes*, *Clostridium perfringens* and *Corynebacterium jeikeium*. Its activity against enterococci is not affected by the presence of *vanA*-, *vanB*- and *vanC*-encoded vancomycin resistance, or by aminoglycoside resistance. Its activity against pneumococci and viridans streptococci is not affected by the presence of intermediate- or high-level penicillin resistance. In addition, the presence of methicillin resistance does not affect the activity of oritavancin against *Staph. aureus* or coagulase-negative staphylococci. Oritavancin is inactive against Gram-negative aerobes and anaerobes [62].

Oritavancin is active in a number of animal models of infection, including a central venous catheter-associated infection model in rats (vancomycin-resistant *E. faecalis*), a rabbit endocarditis model (MRSA, vancomycin-sensitive and vancomycin-resistant *E. faecalis*), a neutropenic mouse thigh model (*Staph. aureus* ATCC 13709), and a rabbit meningitis model (penicillin-sensitive *Strep. pneumoniae*) [63–68].

In a phase 2 clinical trial involving complicated skin and skin-structure infections, oritavancin was not inferior to vancomycin and cephalexin, but had a shorter mean duration of activity [69]. In a phase 2 open-label randomised trial comparing oritavancin (5–10 mg/kg once-daily for 10–14 days) with vancomycin (15 mg/kg twice-daily) and a β -lactam for 10–14 days in patients with *Staph. aureus*-associated bacteraemia, oritavancin was as effective as the comparators, with higher clinical and bacteriological success in the cohort receiving 10 mg/kg, and with no evidence of increased side-effects [70]. The exceptionally long terminal half-life suggests the existence of storage sites within the organism. Studies using cultured macrophages indicate that the drug accumulates slowly (by an endocytic process), but importantly, in the lysosomes, from which efflux is extremely slow [71]. The clinical importance of this property is currently unknown.

Telavancin

Telavancin (TD-6424) is a semi-synthetic derivative of vancomycin, possessing a hydrophobic side-chain on the vancosamine sugar (decylaminoethyl) and a (phosphonomethyl) aminoethyl substituent on the cyclic peptidic core [72]. Pharmacological studies suggest that the enhanced activity of telavancin against *Strep. pneumoniae*, *Staph. aureus* (to a lesser extent) and staphylococci and enterococci harbouring the *vanA* gene cluster results from a complex mechanism of action, which, on the basis of data obtained with close analogues, involves a perturbation of lipid synthesis and possible membrane disruption [73].

Telavancin is active *in vitro* against all Gram-positive pathogens, including *vanA*-positive enterococci. As with oritavancin, extreme potency is observed for streptococci, particularly *Strep. pneumoniae* [74]. Telavancin is active in an *in-vitro* biofilm model, whereas vancomycin and a number of other antibiotics were much less effective [75]. Telavancin is highly effective in animal models of relevant infections [76–78], and tissue penetration of telavancin is high after intravenous administration in healthy subjects [79]. Telavancin has proven to be effective and safe for patients with skin and soft-tissue infections, with cure rates of up to 96%, compared with 90% for standard therapy. Concerns related to QTc elevation will be clarified following further studies [80,81].

Dalbavancin

This agent is a semi-synthetic derivative of the teicoplanin-related glycopeptide A40925, modified with an amide appendage at the C-terminus and an alteration of the hydrophobic acylglucosamine substituent. Like teicoplanin, dalbavancin is active against *vanB*-positive enterococci, as well as staphylococci and other important species [73]. The pharmacokinetic profile of dalbavancin is characterised by a long half-life of *c.* 7 days. This allows a once-weekly regimen for the treatment of infections caused by Gram-positive bacteria. This weekly regimen may offer certain advantages, e.g., improved patient compliance and reduced use of resources compared with antimicrobial agents administered more frequently [82]. Dalbavancin has been shown to be as effective as linezolid twice-daily for the treatment of compli-

cated skin and skin-structure infections [83], and is more effective than most anti-Gram-positive agents, both *in vitro* and *in vivo* [84,85].

In addition to complicated skin and skin-structure infections, dalbavancin has been used to treat catheter-related bloodstream infection. In a randomised controlled open-label multicentre phase 2 trial that involved 75 adult patients with catheter-related bloodstream infection, the results were overwhelmingly in favour of dalbavancin, albeit with limitations [86]. To date, adverse events have been mild and limited, with the most common being pyrexia, headache, nausea, oral candidiasis, diarrhoea and constipation. Consequently, dalbavancin appears to be a promising antimicrobial agent for the treatment of infections caused by Gram-positive bacteria.

CEFTOBIPROLE

Ceftobiprole is a novel broad-spectrum β -lactamase-stable parenteral cephalosporin with strong affinity for the penicillin-binding proteins PBP2a and PBP2x, responsible for resistance in staphylococci and pneumococci, respectively. Ceftobiprole can also bind to relevant penicillin-binding proteins of resistant Gram-positive and Gram-negative bacteria, and appears to have a low ability to select for resistance [87,88]. *In-vivo* screening models suggest good activity of ceftobiprole against Gram-positive and Gram-negative bacteria. Several animal models have been used in the evaluation of ceftobiprole, including mouse sepsis, abscess and pneumonia models, rat endocarditis and tissue cage models, and a rabbit endocarditis model [89,90]. These models suggest that ceftobiprole should have clinical efficacy as an empirical treatment for severe clinical infections.

Several phase 3 studies of ceftobiprole for the treatment of complicated skin and skin-structure infections have been performed. One study compared intravenous ceftobiprole (500 mg every 12 h) with intravenous vancomycin (1 g every 12 h) in patients with complicated skin and skin-structure infections caused by Gram-positive bacteria. Staphylococci were the predominant pathogens, and >25% of the microbiologically evaluable patients had infections caused by MRSA. In the clinically evaluable population, efficacy and adverse events were comparable between treatment arms. Additional clinical trials involving

complicated skin and skin-structure infections and pneumonia patients are underway to evaluate ceftobiprole as a treatment for infections caused by both Gram-positive and Gram-negative bacteria. The broad-spectrum activity of ceftobiprole may allow it to be used as monotherapy for serious nosocomial infections for which combination therapy would otherwise be required [91,92].

CONCLUSIONS

Linezolid, quinupristin-dalfopristin, daptomycin, tigecycline, new glycopeptides and ceftobiprole are the most important novel agents that are currently being considered as alternatives for the treatment of infections caused by Gram-positive bacteria. Doripenem, iclaprim (a recent example of a novel diamonipyrinidine), ranbezolide (a novel oxazolidinone) and ceftaroline (a novel cephalosporin with impressive anti-MRSA and anti-pneumococcal activity) are other agents that are currently under investigation [93–96].

Changing patterns of resistance have compounded and exacerbated the need for new antimicrobial agents. Each of the above compounds has its own unique advantages and disadvantages, so each agent should be used cautiously when conventional treatment fails. The appropriate indications and cost-effectiveness of these molecules will determine future treatment options. Until then, prudent use of existing antibiotics, with strict reinforcement of infection control precautions, should continue to be the rule in the treatment of Gram-positive infections.

ACKNOWLEDGEMENTS

S. Unal is the Turkish coordinator of the linezolid nosocomial pneumoniae study and the TEST tigecycline study group. He has also received travel grants to attend conferences from Pfizer. The authors declare that they have no other conflicts of interest in relation to this manuscript.

REFERENCES

- Johnson AP, Livermore DM, Tillotson GS. Antimicrobial susceptibility of Gram-positive bacteria: what's current, what's anticipated? *J Hosp Infect* 2001; **49** (suppl A): S3–S11.
- Leclercq R, Derlot E, Duval J, Courvalin P. Plasmid-mediated resistance to vancomycin and teicoplanin in *Enterococcus faecium*. *N Engl J Med* 1988; **319**: 157–160.
- Uttley AH, Collins CH, Naidoo J, George RC. Vancomycin-resistant enterococci. *Lancet* 1988; **i**: 57–58.
- Aksoy DY, Tanriover MD, Unal S. Antimicrobial resistance: preventable or inevitable. Problem of the era from two perspectives. In: Gould I, van der Meer J, eds. *Antibiotic policies: fighting resistance*, 1st edn. New York: Springer, 2007: 113–133.
- Anonymous. CDC National Nosocomial Infections Surveillance (NNIS) system report, data summary from January 1990 to May 1999. *Am J Infect Control* 1999; **27**: 520–532.
- Kobayashi K, Rao M, Keis S, Reiney FA, Smith JM, Cook GM. Enterococci with reduced susceptibility to vancomycin in New Zealand. *J Antimicrob Chemother* 2000; **46**: 405–410.
- Appelbaum PC. MRSA—the tip of the iceberg. *Clin Microbiol Infect* 2006; **12** (suppl 2): 3–10.
- Chen CY, Tang JL, Hsueh PR *et al.* Trends and antimicrobial resistance of pathogens causing bloodstream infections among febrile neutropenic adults with hematological malignancy. *J Formos Med Assoc* 2004; **103**: 526–532.
- The European Committee on Antimicrobial Susceptibility Testing (EUCAST) Steering Committee. EUCAST technical note on linezolid. *Clin Microbiol Infect* 2006; **12**: 1243–1245.
- Deshpande LM, Fritsche TR, Moet GJ, Biedenbach DJ, Jones RN. Antimicrobial resistance and molecular epidemiology of vancomycin-resistant enterococci from North America and Europe: a report from the SENTRY antimicrobial surveillance program. *Diagn Microbiol Infect Dis* 2007; **58**: 163–170.
- Gonzales RD, Schreckenberger PC, Graham MB, Kelkar S, DenBesten K, Quinn JP. Infections due to vancomycin-resistant *Enterococcus faecium* resistant to linezolid. *Lancet* 2001; **357**: 1179.
- Toh SM, Xiong L, Arias CA *et al.* Acquisition of a natural resistance gene renders a clinical strain of methicillin-resistant *Staphylococcus aureus* resistant to the synthetic antibiotic linezolid. *Mol Microbiol* 2007; **64**: 1506–1514.
- Ross JE, Fritsche TR, Sader HS, Jones RN. Oxazolidinone susceptibility patterns for 2005: international report from Zyvox Annual Appraisal of Potency and Spectrum study. *Int J Antimicrob Agents* 2007; **29**: 295–301.
- Mylona E, Fanourgiakis P, Vryonis E *et al.* Linezolid-based therapy in *Staphylococcus epidermidis* endocarditis. *Int J Antimicrob Agents* 2007; **29**: 597–598.
- Munoz P, Rodrigues-Creixems M, Moreno M *et al.* Linezolid therapy for infective endocarditis. *Clin Microbiol Infect* 2007; **13**: 211–215.
- Moitra RK, Auckland C, Cawley MI, Jones G, Cooper C. Systemic nocardiosis in a splenectomized patient with systemic lupus erythematosus. Successful treatment using linezolid. *J Clin Rheumatol* 2003; **9**: 47–50.
- Sood R, Bhadauriya T, Rao M *et al.* Antimycobacterial activities of oxazolidinones: a review. *Infect Disord Drug Targets* 2006; **6**: 343–354.
- Ntziora F, Falagas ME. Linezolid for the treatment of patients with central nervous system infection. *Ann Pharmacother* 2007; **41**: 296–308.
- Falagas ME, Siempos II, Papagelopoulos PJ, Vardakas KZ. Linezolid for the treatment of adults with bone and joint infections. *Int J Antimicrob Agents* 2007; **29**: 233–239.
- Kutscha-Lissberg F, Hebler U, Muhr G, Koller M. Linezolid penetration into bone and joint tissues infected with methicillin-resistant staphylococci. *Antimicrob Agents Chemother* 2003; **47**: 3964–3966.

21. Wunderink RG, Rello J, Cammarata SK, Croos-Dabrera RV, Kollef MH. Linezolid vs vancomycin, analysis of two double blind studies of patients with methicillin-resistant *Staphylococcus aureus* nosocomial pneumonia. *Chest* 2003; **124**: 1789–1797.
22. Wilson SE. Clinical trial results with linezolid, an oxazolidinone, in the treatment of soft tissue and post-operative Gram positive infections. *Surg Infect (Larchmt)* 2001; **2**: 25–35.
23. Hau T. Efficacy and safety of linezolid in the treatment of skin and soft tissue infections. *Eur J Clin Microbiol Infect Dis* 2002; **21**: 491–498.
24. Maclayton DO, Hall RG. Pharmacologic treatment options for nosocomial pneumonia involving methicillin-resistant *Staphylococcus aureus*. *Ann Pharmacother* 2007; **41**: 235–244.
25. Ruiz ME, Guerrero IC, Tuazon CU. Endocarditis caused by methicillin-resistant *Staphylococcus aureus*: treatment failure with linezolid. *Clin Infect Dis* 2002; **35**: 1018–1020.
26. Schweiger ES, Weinberg JM. Novel antibacterial agents for skin and skin structure infections. *J Am Acad Dermatol* 2004; **50**: 331–340.
27. Bishop E, Melvani S, Howden BP, Charles PG, Grayson ML. Good clinical outcomes but high rates of adverse reactions during linezolid therapy for serious infections: a proposed protocol for monitoring therapy in complex patients. *Antimicrob Agents Chemother* 2006; **50**: 1599–1602.
28. Bryson HM, Spencer CM. Quinupristin–dalfopristin: antibacterial activity, pharmacokinetic profile, therapeutic trials and tolerability and current status. *Drugs* 1996; **52**: 406–415.
29. Cocito C, Di Giambattista M, Nyssen E, Vannuffel P. Inhibition of protein synthesis by streptogramins and related antibiotics. *J Antimicrob Chemother* 1997; **39** (suppl A): 7–13.
30. Johnson AP, Livermore DM. Quinupristin/dalfopristin: a new addition to antimicrobial arsenal. *Lancet* 1999; **354**: 2012–2013.
31. Nichols RL, Graham DR, Barriere SL *et al.* Treatment of hospitalized patients with complicated gram-positive skin and skin structure infections: two randomized, multicentre studies of quinupristin–dalfopristin versus cefazolin, oxacillin or vancomycin. Synercid Skin and Skin Structure Infection Group. *J Antimicrob Chemother* 1999; **44**: 263–273.
32. Eliopoulos G. Antimicrobial agents for treatment of serious infections caused by resistant *Staphylococcus aureus* and enterococci. *Eur J Clin Microbiol Infect Dis* 2005; **24**: 826–831.
33. Moellering RC, Linden PK, Reinhardt J, Blumberg EA, Bompert F, Talbot GH. The efficacy and safety of quinupristin/dalfopristin for the treatment of infections caused by vancomycin-resistant *Enterococcus faecium*. Synercid Emergency-Use Study Group. *J Antimicrob Chemother* 1999; **44**: 251–261.
34. Stamatakis MK, Richards JG. Interaction between quinupristin/dalfopristin and cyclosporine. *Ann Pharmacother* 1997; **31**: 576–578.
35. Torres-Viera C, Dembry LM. Approaches to vancomycin-resistant enterococci. *Curr Opin Infect Dis* 2004; **17**: 541–547.
36. Ginsburg I. The role of bacteriolysis in the pathophysiology of inflammation, infection and post infectious sequelae. *APMIS* 2002; **110**: 753–770.
37. Akins RL, Rybak MJ. Bactericidal activities of two daptomycin regimens against clinical strains of glycopeptide intermediate-resistant *Staphylococcus aureus*, vancomycin-resistant *Enterococcus faecium*, and methicillin-resistant *Staphylococcus aureus* isolates in an *in vitro* pharmacodynamic model with simulated endocardial vegetations. *Antimicrob Agents Chemother* 2001; **45**: 454–459.
38. Rybak MJ, Hershberger E, Moldovan T, Grucz RG. *In vitro* activities of daptomycin, vancomycin, linezolid and quinupristin/dalfopristin against staphylococci and enterococci including vancomycin-intermediate and resistant strains. *Antimicrob Agents Chemother* 2000; **44**: 1062–1066.
39. Cha R, Grucz RG, Rybak MJ. Daptomycin dose–effect relationship against resistant gram-positive organisms. *Antimicrob Agents Chemother* 2003; **47**: 1598–1603.
40. Arbeit RD, Maki D, Tally FP *et al.* The safety and efficacy of daptomycin for the treatment of complicated skin and skin-structure infections. *Clin Infect Dis* 2004; **38**: 1673–1661.
41. Martone WJ, Lamp KC. Efficacy of daptomycin in complicated skin and skin-structure infections due to methicillin-sensitive and -resistant *Staphylococcus aureus*: results from the CORE registry. *Curr Med Res Opin* 2006; **22**: 2337–2343.
42. Fowler VG, Boucher HW, Corey GR *et al.* S. aureus Endocarditis and Bacteremia Study Group. Daptomycin versus standard therapy for bacteremia and endocarditis caused by *Staphylococcus aureus*. *N Engl J Med* 2006; **355**: 653–665.
43. Segreti JA, Crank CW, Finney MS. Daptomycin for the treatment of gram-positive bacteremia and infective endocarditis. A retrospective case series of 31 patients. *Pharmacotherapy* 2006; **26**: 347–352.
44. Vouillamoz J, Moreillon P, Giddey M, Entenza JM. Efficacy of daptomycin in the treatment of experimental endocarditis due to susceptible and multi-drug resistant enterococci. *J Antimicrob Chemother* 2006; **58**: 1208–1214.
45. Silverman JA, Mortin LI, Vanpraagh AD, Li T, Alder J. Inhibition of daptomycin by pulmonary surfactant: *in vitro* modeling and clinical impact. *J Infect Dis* 2005; **191**: 2149–2152.
46. Falagas ME, Giannopoulou KP, Ntziora F, Vardakas KZ. Daptomycin for endocarditis and/or bacteraemia: a systematic review of the experimental and clinical evidence. *J Antimicrob Chemother* 2007; **60**: 7–19.
47. Mangili A, Bica I, Snyderman DR, Hamer DH. Daptomycin-resistant methicillin-resistant *Staphylococcus aureus* bacteremia. *Clin Infect Dis* 2005; **40**: 1058–1060.
48. Skiest DJ. Treatment failure resulting from resistance of *Staphylococcus aureus* to daptomycin. *J Clin Microbiol* 2006; **44**: 655–656.
49. Hoban DJ, Bouchillon SK, Johnson BM *et al.* *In vitro* activities of tigecycline against 6792 gram-negative and gram-positive clinical isolates from the global Tigecycline Evaluation and Surveillance Trial (TEST Program, 2004). *Diagn Microbiol Infect Dis* 2005; **52**: 215–227.
50. Pankey GA. Tigecycline. *J Antimicrob Chemother* 2005; **56**: 470–480.
51. Projan SJ. Preclinical pharmacology of GAR-936, a novel glycylcycline antibacterial agent. *Pharmacotherapy* 2000; **20**: 219S–223S.
52. Sum PE, Petersen P. Synthesis and structure–activity relationship of novel glycylcycline derivatives leading to the discovery of GAR-936. *Bioorg Med Chem Lett* 1999; **9**: 1459–1462.

53. Al-Tatari H, Abdel-Haq N, Chearskul P, Asmar B. Antibiotics for treatment of resistant Gram-positive coccal infections. *Ind J Pediatr* 2006; **73**: 323–334.
54. Sacchidanand S, Penn RL, Embil JM *et al*. Efficacy and safety of tigecycline monotherapy compared with vancomycin plus aztreonam in patients with complicated skin and skin structure infections: results from a phase 3, randomized, double-blind trial. *Int J Infect Dis* 2005; **9**: 251–261.
55. Ellis-Grosse EJ, Babichak T, Dartois N *et al*. The efficacy and safety of tigecycline in the treatment of skin and skin structure infections: results of 2 double-blind phase 3 comparison studies with vancomycin–aztreonam. *Clin Infect Dis* 2005; **41** (suppl 5): S341–S353.
56. Babinchak T, Ellis-Grosse E, Dartois N *et al*. The efficacy and safety of tigecycline for the treatment of complicated intra-abdominal infections: analysis of pooled clinical trial data. *Clin Infect Dis* 2005; **41** (suppl 5): S354–S367.
57. Oliva ME, Rekha A, Yellin A *et al*. A multicenter trial of the efficacy and safety of tigecycline versus imipenem/cilastatin in patients with complicated intra-abdominal infections. *BMC Infect Dis* 2005; **5**: 88.
58. Stein GE, Craig WA. Tigecycline: a critical analysis. *Clin Infect Dis* 2006; **43**: 518–524.
59. Cooper RD, Snyder NJ, Zweifel MJ *et al*. Reductive alkylation of glycopeptide antibiotics: synthesis and antibacterial activity. *J Antibiot* 1996; **49**: 575–581.
60. Malabarba A, Ciabatti R. Glycopeptide derivatives. *Curr Med Chem* 2001; **8**: 1759–1773.
61. Allen NE, Nicas TI. Mechanism of action of oritavancin and related glycopeptide antibiotics. *FEMS Microbiol Rev* 2003; **26**: 511–532.
62. Guay DR. Oritavancin and tigecycline: investigational antimicrobials for multi-drug resistant bacteria. *Pharmacotherapy* 2004; **24**: 58–68.
63. Saleh-Mghir A, Lefort A, Petegnief Y *et al*. Activity and diffusion of LY333328 in experimental endocarditis due to vancomycin-resistant *Enterococcus faecalis*. *Antimicrob Agents Chemother* 1999; **43**: 115–120.
64. Lefort A, Saleh-Mghir A, Garry L, Carbon C, Fantin B. Activity of LY333328 combined with gentamicin *in vitro* and in rabbit experimental endocarditis due to vancomycin-susceptible or -resistant *Enterococcus faecalis*. *Antimicrob Agents Chemother* 2000; **44**: 3017–3021.
65. Rupp ME, Fey PD, Longo GM. Effect of LY333328 against vancomycin-resistant *Enterococcus faecium* in a rat central venous catheter-associated infection model. *J Antimicrob Chemother* 2001; **47**: 705–707.
66. Kaatz GW, Seo SM, Aeschlimann JR, Houlihan HH, Mercier RC, Rybak MJ. Efficacy of LY333328 against experimental methicillin-resistant *Staphylococcus aureus* endocarditis. *Antimicrob Agents Chemother* 1998; **42**: 981–983.
67. Gerber J, Smirnov A, Wellmer A *et al*. Activity of LY333328 in experimental meningitis caused by a *Streptococcus pneumoniae* strain susceptible to penicillin. *Antimicrob Agents Chemother* 2001; **45**: 2169–2172.
68. Boylan CJ, Campanale K, Iverson PW, Phillips DL, Zeckel ML, Parr TR. Pharmacodynamics of oritavancin (LY333328) in a neutropenic-mouse thigh model of *Staphylococcus aureus* infection. *Antimicrob Agents Chemother* 2003; **47**: 1700–1706.
69. Mercier RC, Hrebickova L. Oritavancin: a new avenue for resistant Gram-positive bacteria. *Expert Rev Anti-Infect Ther* 2005; **3**: 325–332.
70. Loutit JS, O'Riordan W, San Juan J *et al*. Phase 2 trial comparing four regimens of oritavancin vs. comparator in the treatment of patients with *S. aureus* bacteremia. *Clin Microbiol Infect* 2004; **10** (suppl 3): 122.
71. Van Bambeke F, Carryn S, Seral C *et al*. Cellular pharmacokinetics and pharmacodynamics of oritavancin (LY333328) glycopeptide in a model of J774 mouse macrophages. *Antimicrob Agents Chemother* 2004; **48**: 2853–2860.
72. Judice JK, Pace JL. Semi-synthetic glycopeptide antibacterials. *Bioorg Med Chem Lett* 2003; **13**: 4165–4168.
73. Van Bambeke F. Glycopeptides in clinical development: pharmacological profile and clinical perspectives. *Curr Opin Pharmacol* 2004; **4**: 471–478.
74. Pace JL, Yang G. Glycopeptides: update on an old successful antibiotic class. *Biochem Pharmacol* 2006; **71**: 968–980.
75. Gander S, Kinnaird A, Finch R. Telavancin: *in vitro* activity against staphylococci in a biofilm model. *J Antimicrob Chemother* 2005; **56**: 337–343.
76. Reyes N, Skinner R, Benton BM *et al*. Efficacy of telavancin in a murine model of bacteraemia induced by methicillin-resistant *Staphylococcus aureus*. *J Antimicrob Chemother* 2006; **58**: 462–465.
77. Stucki A, Gerber P, Acosta F, Cottagnoud M, Cottagnoud P. Efficacy of telavancin against penicillin-resistant pneumococci and *Staphylococcus aureus* in a rabbit meningitis model and determination of kinetic parameters. *Antimicrob Agents Chemother* 2006; **50**: 770–773.
78. Madrigal AG, Basuino L, Chambers HF. Efficacy of telavancin in a rabbit model of aortic valve endocarditis due to methicillin-resistant *Staphylococcus aureus* or vancomycin-intermediate *Staphylococcus aureus*. *Antimicrob Agents Chemother* 2005; **49**: 3163–3165.
79. Sun HK, Duchin K, Nightingale CH, Shaw JP, Seroogy J, Nicolau DP. Tissue penetration of telavancin after intravenous administration in healthy subjects. *Antimicrob Agents Chemother* 2006; **50**: 788–790.
80. Stryjewski ME, Chu VH, O'Riordan WD *et al*. Telavancin versus standard therapy for treatment of complicated skin and skin structure infections caused by gram positive bacteria; FAST 2 study. *Antimicrob Agents Chemother* 2006; **50**: 862–867.
81. Stryjewski ME, O'Riordan WD, Lau WK *et al*. Telavancin versus standard therapy for treatment of complicated skin and skin structure infections due to gram positive bacteria. *Clin Infect Dis* 2005; **40**: 1601–1607.
82. Dorr MB, Jabes D, Cavaleri M *et al*. Human pharmacokinetics and rationale for once-weekly dosing of dalbavancin, a semisynthetic glycopeptide. *J Antimicrob Chemother* 2005; **55** (suppl 2): 25–30.
83. Jauregui LE, Babazadeh S, Seltzer E *et al*. Randomized, double-blind comparison of once weekly dalbavancin versus twice-daily linezolid therapy for the treatment of complicated skin and skin structure infections. *Clin Infect Dis* 2005; **41**: 1407–1415.
84. Lefort A, Pavie J, Garry L, Chau F, Fantin B. Activities of dalbavancin *in vitro* and in a rabbit model of experimental endocarditis due to *Staphylococcus aureus* with or without reduced susceptibility to vancomycin and teicoplanin. *Antimicrob Agents Chemother* 2004; **48**: 1061–1064.
85. Malabarba A, Goldstein BP. Origin, structure and activity *in vitro* and *in vivo* of dalbavancin. *J Antimicrob Chemother* 2005; **55** (suppl 2): 15–20.

86. Raad I, Darouiche R, Vazquez J *et al.* Efficacy and safety of weekly dalbavancin therapy for catheter-related bloodstream infections caused by Gram-positive pathogens. *Clin Infect Dis* 2005; **40**: 374–380.
87. Hebeisen P, Heinze-Krauss I, Angehrn P, Hohl P, Page MG, Then RL. *In vitro* and *in vivo* properties of Ro-63-9141, a novel broad-spectrum cephalosporin with activity against methicillin-resistant staphylococci. *Antimicrob Agents Chemother* 2001; **45**: 825–836.
88. Bogdanovich T, Ednie LM, Shapiro S, Appelbaum PC. Antistaphylococcal activity of ceftobiprole: a new broad spectrum cephalosporin. *Antimicrob Agents Chemother* 2005; **49**: 4210–4219.
89. Chambers HF. Evaluation of ceftobiprole in a rabbit model of aortic valve endocarditis due to methicillin-resistant and vancomycin intermediate *Staphylococcus aureus*. *Antimicrob Agents Chemother* 2005; **49**: 884–888.
90. Vaudaux P, Gjinovci A, Bento M, Li D, Schrenzel J, Lew DP. Intensive therapy with ceftobiprole medocaril of experimental foreign-body infection by methicillin-resistant *Staphylococcus aureus*. *Antimicrob Agents Chemother* 2005; **49**: 3789–3793.
91. Noel GJ. Clinical profile of ceftobiprole, a novel beta-lactam antibiotic. *Clin Microbiol Infect* 2007; **13** (suppl 2): 25–29.
92. Chambers HF. Ceftobiprole: *in vivo* profile of a bactericidal cephalosporin. *Clin Microbiol Infect* 2006; **12** (suppl 2): 17–22.
93. Mushtaq S, Warner M, Ge Y, Kaniga K, Livermore DM. *In vitro* activity of ceftaroline (PPI-0903M, T-91825) against bacteria with defined resistance mechanisms and phenotypes. *J Antimicrob Chemother* 2007; **60**: 300–311.
94. Poulakou G, Giamarellou H. Investigational treatments for postoperative surgical site infections. *Expert Opin Investig Drugs* 2007; **16**: 137–155.
95. Hawser S, Lociuro S, Islam K. Dihydrofolate reductase inhibitors as antibacterial agents. *Biochem Pharmacol* 2006; **71**: 941–948.
96. Mathur T, Bhateja P, Pandya M, Fatma T, Rattan A. *In vitro* activity of RBx 7644 (ranbezolid) on biofilm producing bacteria. *Int J Antimicrob Agents* 2004; **24**: 369–373.